## STREPGUARD® WITH HAVLOGEN®\*

## Intervet

Streptococcus Equi Bacterial Extract

An adjuvanted, concentrated, enzyme extract of Streptococcus equi. \*+

Recommended for use in healthy horses as an aid in the prevention of strangles disease due to *Streptococcus equi* infection.

**DOSE:** Inject 1 mL intramuscularly using aseptic techniques. Repeat in 3-4 weeks. A 1 mL dose should be administered annually or prior to expected exposure.

Withdrawal Time: Do not vaccinate within 21 days of slaughter.

**CAUTION:** Store at 35° to 45°F (2° to 7°C). Shake well before using. Use entire contents when first opened. Do not vaccinate within 21 days before slaughter. Certain hypersensitive individuals may demonstrate local or generalized reactions, sometimes severe, following exposure to streptococcal proteins. To minimize reactions, free exercise is recommended after injection. Anaphylactoid reactions may occur.

**ANTIDOTE:** Epinephrine. Contains thimerosal as preservative.

\*Adjuvant - U.S. Patent Nos. 3,790,665 and 3,919,411.

+U.S. Patent No. 4,582,798.

FOR USE IN ANIMALS ONLY

U.S. Vet. License No. 286

**SUPPLIED:** 

Code 4666 10

10 Doses

Ten 1 mL Syringes with Separate

77046660, R.5

Sterile Needle

Code 4667

10 Doses

10 mL

76046670, R.5

Disclaimer: Every effort has been made to ensure the accuracy of the information published. However, it remains the responsibility of the readers to familiarize themselves with the product information contained on the product label or package insert. Compendium Code No.: 08391195